



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2010

**A multiplex-system to target 16 male-specific and 15 autosomal genetic
markers for orang-utans (genus: Pongo)**

Nietlisbach, Pirmin ; Nater, Alexander ; Greminger, Maja P ; Arora, Natasha ; Krützen, Michael

DOI: <https://doi.org/10.1007/s12686-010-9278-2>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-71190>

Journal Article

Published Version

Originally published at:

Nietlisbach, Pirmin; Nater, Alexander; Greminger, Maja P; Arora, Natasha; Krützen, Michael (2010).
A multiplex-system to target 16 male-specific and 15 autosomal genetic markers for orang-utans (genus:
Pongo). Conservation Genetics Resources, 2(S1):153-158.

DOI: <https://doi.org/10.1007/s12686-010-9278-2>

A multiplex-system to target 16 male-specific and 15 autosomal genetic markers for orang-utans (genus: *Pongo*)

Pirmin Nietlisbach · Alexander Nater ·
Maja P. Greminger · Natasha Arora ·
Michael Krützen

Received: 30 June 2010 / Accepted: 6 July 2010 / Published online: 25 July 2010
© Springer Science+Business Media B.V. 2010

Abstract Genetic studies of dispersal on local spatial and short temporal scales require a large number of autosomal microsatellites. However, the study of dispersal over large spatial scales and the resolution of deep evolutionary histories require marker systems that are preferentially inherited through the male or female line. Addressing such questions in endangered orang-utans (genus: *Pongo*) bears significant relevance to species conservation, as habitat destruction and fragmentation pose a significant threat to the whole genus. Here, we report 16 male-specific markers (nine human-derived microsatellites, six single nucleotide and one insertion-deletion polymorphisms), and 15 novel *Pongo*-derived autosomal microsatellite loci. All 31 markers can be amplified in four multiplex polymerase chain reactions even in DNA derived from faecal material. The markers can be applied to studying a wide range of important questions in this genus, such as conservation genetics, social structure, phylogeny and phylogeography.

Keywords *Pongo* spp. ·

Single nucleotide polymorphisms · Microsatellites ·
Y chromosome · SNP typing · Non-invasive samples

The endangered orang-utans occur on the islands of Borneo (*Pongo pygmaeus*; about 50,000 animals) and Sumatra (*P. abelii*; about 6,500 animals), where they have undergone a recent dramatic decline in population size (Goossens et al. 2006; Wich et al. 2008). This has been mostly attributed to habitat loss, leading to heavily fragmented populations of often only a few hundred individuals (Wich et al. 2008). Therefore, it is essential to maintain genetic diversity, which has been linked to population fitness (e.g. Reed and Frankham 2003). This can be achieved by maintaining corridors between fragmented populations, allowing animals to follow natural dispersal patterns (Gilbert-Norton et al. 2010).

Studying natural dispersal in wild orang-utans pose significant challenges. Behavioural observations suggested higher male than female dispersal (Delgado and van Schaik 2000), although this has not been fully confirmed by previous genetic studies (Utami et al. 2002; Goossens et al. 2005), where patterns of direct dispersal were investigated using autosomal microsatellite markers. However, direct inferences from autosomal markers are limited to the timescale of a few generations and geographically small areas, as sexual recombination will break down sex-specific information (Goudet et al. 2002). Sex-biased dispersal over larger time and spatial scales can be investigated by contrasting genetic information obtained from markers inherited through either the male or female lineage (Handley and Perrin 2007).

In orang-utans, maternally transmitted mitochondrial DNA markers are widely available (e.g. Warren et al. 2001), but markers on the male-specific region of the Y chromosome have not yet been applied. Here, we report 16 male-specific markers for the application in the genus *Pongo*. Nine of these markers are human-derived microsatellite loci, six are single nucleotide polymorphisms (SNPs) and one is an

Electronic supplementary material The online version of this article (doi:10.1007/s12686-010-9278-2) contains supplementary material, which is available to authorized users.

P. Nietlisbach · A. Nater · M. P. Greminger · N. Arora ·
M. Krützen (✉)
Evolutionary Genetics Group, Anthropological Institute
and Museum, University of Zurich, Winterthurerstrasse 190,
8057 Zurich, Switzerland
e-mail: michael.krutzen@aim.uzh.ch

Table 1 Primers for male-specific and autosomal markers in orang-utans

Locus	Primer sequence 5'–3'	PC	T _A	Polymorphism	Overall		Suaiq (Sumatra)			Tuanan (Borneo)		
					N	N _A	N	N _A	H _O	N	N _A	H _O
DYS630	PET-AGCAAGACTCCACCTCAAAAAGA*	0.15	63	AGAA	172	11	14	3	0.66	21	4	0.69
	gtttGCTGTGAGTTCATATAATTTCTTCC	0.20		indel	172	2	14	1	0	21	1	0
DYS587	6FAM-AAAAATTACCTTCTTTGGAAAAGTAGTATT	0.30	63	ATACA	166	8	14	1	0	19	1	0
	gtttGTTATTCTGAGCAGGGTTTCTAAG	0.40										
DYS532	NED-AGCAGGATTCCTCTAAAAATATCA	0.10	63	compound, main motif	171	3	14	1	0	21	2	0.09
	gTTTCTCCCTCCCTCCCTCTC	0.14		(CTTT)								
DYS577	6FAM-CCACTAAGCCCATGCATATTATT	0.30	63	GAAT	171	2	14	1	0	19	1	0
	gtttGAGAGGTTGAGGCTGCAGTAAG	0.40		C/G	171	2	14	2	0.13	19	1	0
	gtttGAGAGGTTGAGGCTG CAGTAAC	0.40										
DYS645	6FAM-GTACTAATTTTATTCTTATGGCGTAGA	0.15	63	GTTTT	173	2	14	1	0	21	1	0
	gtttACACATGGCACCTGACACTG	0.20										
Y6C2	6FAM-CTTCTCTCTCTCTCTCTCTCTCTCTCT	0.10	63	TTC	172	2	14	1	0	20	1	0
	gtttCAATAGTTTGGGAAATAAGACAAATG	0.14										
DBY13	6FAM-GGAAACTAAAAATATGACATTGTAAATTG	0.30	63	C/G	168	2	14	1	0	20	1	0
	gtttAATTTTATTATGTGATGCATACAGC	0.40										
	gtttGATTTTATTTTATTGTGATGCATACAGG	0.40										
DYS510	PET-GAAAGATAGATCAACAAGGTAGAAACAA	0.30	64	GATA	169	6	12	4	0.6	21	2	0.44
	gtttCATCCATCCATCCATCCATCT	0.40										
DYS561	6FAM-CCTGATGCCATCTGAAAAATTAA	0.30	64	TAGA	168	5	14	1	0	20	3	0.52
	gtttACAACTGCCTCCAGCTTAGG	0.40										
DYS556	6FAM-TTACAAAACTAACATAAAGACCAACACAG	0.30	64	TAAA	172	3	14	1	0	21	2	0.41
	gtttGAAGCATTTGAGTATAGTATAAAGTTGGT	0.40										
DYS630	PET-AGCAAGACTCCACCTCAAAAAGA*	0.15	64	A/G	171	2	14	1	0	21	1	0
	gtttTGAGTTCCATAAATTTCTCTCTTCC	0.20										
	gtttGTGAGTTCCATAAATTTCTCTCTTCT	0.20										
SMCY12_26	6FAM-AAGGGTCACACAGAAATACCTTAG	0.15	64	C/G	173	2	14	2	0.13	21	1	0
	gtttGACAGGTGGGGCGTAGTCTC	0.20										
	gtttCAGGTGGGGCGTAGTCTG	0.20										
SMCY12_337	6FAM-GTTACAGGTATACATGCACCTTTTT	0.15	64	A/C	171	2	14	1	0	21	1	0
	gtttGTTGTTGGCTTCTTTACTCTGTCA	0.20										
	gtttGTTGTTGGCTTCTTTACTCTGTCC	0.20										
SMCY14	6FAM-ATGGGAAAAAGATGAGTTCTGA	0.15	64	C/T	173	2	14	1	0	21	2	0.41
	gtttGTCTGGCATCCTAATGCCT	0.20										
	gtttGTCTGGCATCCTAATGCC	0.20										

Table 1 continued

Locus	Primer sequence 5'–3'	PC	T _A	Polymorphism	Overall		Suaiq (Sumatra)				Tuanan (Borneo)			
					N	N _A	N	N _A	H _O	H _E	N	N _A	H _O	H _E
O4_6	PE7-GGCAATGTAACATATCCCTCTGTGT AGCCATGGACCTTGTGAGAAAAG	0.05 0.05	58	GATA			23	4	0.61	0.68	28	3	0.71	0.62
O4_A5	6FAM-ATGGGCCAGAAAACAACTCAGT AGATAAAGGAATGGATAGATGGACAGA	0.15 0.15	58	(GATA)(GATG)			22	4	0.64	0.55	26	6	0.65	0.65
O4_A7	VIC-ATGGGCCCAATCAAAGTCTGTCAAT ACTGGCCCAATCAAAGTCTGT	0.10 0.10	58	GTAG			21	4	0.86	0.72	26	2	0.35	0.29
O4_A8	NED-CACAGGGTCCAAACTCAGATTATTG CCTCCCTCATGTAGTTATCAA	0.20 0.20	58	(GATA)(GATG)			23	3	0.30	0.31	29	1	0	0
O4_B5	VIC-GAGCCCTGATTCGTTTACTGG AGCAAAGGCAGAAAAGTGAATGA	0.20 0.20	58	GATA			22	6	0.86	0.73	28	5	0.50	0.54
O4_B6	6FAM-TGGAGCCTGAATATGTGACTGAAT AATGCCAGGATTTCCCTCTTTT	0.20 0.20	58	(GATA)(GTAG)			20	6	0.65	0.61	26	6	0.46	0.79
O4_B24	6FAM-TCTGAGGTACCCTGTAAACAAAGAAA GAAATCCCAGTACCATATAAATGTCTAT	0.10 0.10	58	GATA			23	3	0.65	0.56	29	1	0.00	0.00
O4_A1	6FAM-CTCCCTTCCTTCCTTTATTCAGTT CAACACTTGGCAGTCACAAATCAG	0.10 0.10	62	GTAG			23	5	0.87	0.73	28	4	0.82	0.75
O4_B3	VIC-TTCCAGAAAGGGCGAGAAAGTT GTTGGACCAACAGTTGTCAATAA	0.10 0.10	62	GACA			22	3	0.59	0.64	26	1	0	0
O4_B17	PE7-GTACGACGGTGCACGAACAATGTA AGCCTGGCTGAAAAGTGGAACTGAG	0.30 0.30	62	GATG			19	3	0.68	0.67	26	6	0.69	0.73
O4_B20	NED-CCTGCATTTTGTCACTCCCTCAACC CTGCCACACCTCCATGGACACAGAT	0.20 0.20	62	GATG			14	1	0	0	24	2	0.33	0.38
O4_C9	6FAM-TGCAGGCCAGGGCTTCTTTCAA CAGTCTCCCAAGGACCCCTACACAG	0.15 0.15	62	GATA			22	5	0.55	0.54	27	4	0.59	0.63
O4_C13	6FAM-CTGGGCACACTGTATATGGGGTAG GTTTGAGACCACTCATGTGCAAAAGACC	0.20 0.20	62	GATA			20	3	0.75	0.56	21	4	0.38	0.59
O4_Chr5	PE7-CAGCAGCTCTGAAATATCTGTCC GTTTGGGTAGAGGAAAAGCAGGTGTAT	0.15 0.15	62	GATA			21	4	0.81	0.70	23	5	0.74	0.74
O4_Chr7	NED-CATCTCTTTATGGCTGACTGTGTAT GTTTGGTCCAAAGACAAATTTGTATGAGT	0.10 0.10	62	GATA			17	11	0.76	0.83	24	15	0.92	0.91

All loci with a Y in the name are Y-linked, all others are autosomal. Summary statistics are given for two study sites and over all sampled orang-utans. For loci DY5630 and DY5577, three and two male-specific markers were typed, respectively (Fig. 1). Loci combined in a single multiplex reaction have the same annealing temperature

Fluorescent labels are shown in italics at the 5' end of the forward primer. PIG-tail bases (Brownstein et al. 1996) are given in lower case

PC primer concentration [μM], T_A annealing temperature, N number of samples, N_A number of different alleles, H_O observed heterozygosity, H_E expected heterozygosity (Nei 1987). * primer used in two PCRs (Fig. 1b)

Amplicon sizes and their relation to repeat numbers are shown in Table S1 in the Online Resources

(Nietlisbach 2009), if used in unison with readily available mtDNA markers.

To clone autosomal microsatellite markers, we extracted genomic DNA from 25 mg of frozen muscle tissue from a Sumatran orang-utan, using the DNeasy Tissue Kit (Qiagen). We digested ten micrograms of the purified DNA with *NheI* and *AluI* (New England Biolabs) and size-selected for fragments between 400 and 1,200 base pairs length. Enrichment, cloning and sequencing were carried out as described in Nater et al. (2008), using only tetra-nucleotide biotinylated probes [(GACA)₇, (GATA)₇, and (GATC)₇]. We sequenced plasmids from 68 positive clones, of which 70% contained a microsatellite repeat. For 25 loci, which contained long uninterrupted repeats, we designed primers and amplified these loci in twelve orang-utans. Levels of polymorphism were qualitatively assessed on high-resolution Spreadex gels (Elchrom Scientific). Based on these results, we fluorescently labelled the forward primers of the 15 most polymorphic markers and combined these 15 loci (GenBank Acc.No. HM804007–HM804021) into two multiplex PCRs (Table 1). Then, we genotyped 29 orang-utans from Borneo and 23 from Sumatra, using DNA extracts from faecal samples with target DNA concentration ranging from 25 to 1,000 pg/μl, strictly following guidelines from Morin et al. (2001). PCRs using the Qiagen PCR Multiplex Kit contained 1 μl template DNA in an 8 μl final volume, with varying primer concentrations and annealing temperatures (Table 1). PCRs included 45 cycles with conditions according to manufacturer's instructions.

If not indicated otherwise, we used standard laboratory techniques at each step. We designed PCR primers with the PrimerSelect software implemented in Lasergene v7 (DNASTAR). PCR amplifications were performed on Veriti 96-well thermal cyclers (Applied Biosystems). Sequencing reactions were carried out using the BigDye Terminator v3.1 on a 3730 DNA Analyzer (both Applied Biosystems) according to manufacturer's instructions, cleaned-up using a MgSO₄ precipitation procedure, followed by resuspending the pellet in 20 μl ddH₂O. For fragment length analysis, PCR products were diluted 20–80 times in ddH₂O. One microlitre of this was added to 9.93 μl HiDi formamide and 0.07 μl of GeneScan 500 LIZ Size Standard (both Applied Biosystems) and denatured for three minutes at 95°C. We ran the samples on a 3730 DNA Analyzer and obtained genotypes using GeneMapper software v4.0 (Applied Biosystems). For the statistical analyses, we used MStools v3.1 add-into Microsoft Excel (Park 2001) and Genepop v4.0 (Rousset 2008).

Fragment length discrepant allele specific PCR used as SNP typing technique proved to be a reliable and cost-efficient strategy to assess SNP variation. The possibility to combine this technique with conventional microsatellite

fragment length analysis makes it a suitable method to include a small number of SNPs to complement an extensive microsatellite analysis. The polymorphic male-specific markers for orang-utans described here promise to be highly useful for population genetic and phylogenetic studies addressing questions about dispersal strategies, phylogeographic patterns, and comparisons with other molecular markers. The autosomal markers can be applied to investigate local dispersal or assess relatedness and paternity. Knowledge about such processes, in particular about natural dispersal strategies, is important for species conservation.

Acknowledgments We thank C. van Schaik, M. van Noordwijk, J. Pamungkas, and D. Perwitasari-Farajallah. We are also indebted to all individuals who helped collecting samples in the field. This study was funded by the Swiss National Science Foundation (31003A-116848 to MK), Messerli Foundation, A.H.-Schultz Stiftung, and Claraz Schenkung. We thank the Indonesian Institute of Sciences (LIPI), the Indonesian State Ministry for Research and Technology (RISTEK), and the Sabah Wildlife Department for granting permission to undertake this research. All sampling and transportation of samples was conducted in accordance with Indonesian, Malaysian and international regulations (CITES).

References

- Brownstein MJ, Carpten JD, Smith JR (1996) Modulation of non-templated nucleotide addition by *Taq* DNA polymerase: primer modifications that facilitate genotyping. *Biotechniques* 20(6): 1004–1010
- Delgado RA, van Schaik CP (2000) The behavioral ecology and conservation of the orang-utan (*Pongo pygmaeus*): a tale of two islands. *Evol Anthropol* 9(5):201–218
- Erler A, Stoneking M, Kayser M (2004) Development of Y-chromosomal microsatellite markers for nonhuman primates. *Mol Ecol* 13(10):2921–2930. doi:10.1111/j.1365-294X.2004.02304.x
- Gilbert-Norton L, Wilson R, Stevens JR, Beard KH (2010) A meta-analytic review of corridor effectiveness. *Conserv Biol* 24(3): 660–668. doi:10.1111/j.1523-1739.2010.01450.x
- Goossens B, Chikhi L, Jalil MF, Ancrenaz M, Lackman-Ancrenaz I, Mohamed M, Andau P, Bruford MW (2005) Patterns of genetic diversity and migration in increasingly fragmented and declining orang-utan (*Pongo pygmaeus*) populations from Sabah, Malaysia. *Mol Ecol* 14(2):441–456. doi:10.1111/j.1365-294X.02421.x
- Goossens B, Chikhi L, Ancrenaz M, Lackman-Ancrenaz I, Andau P, Bruford MW (2006) Genetic signature of anthropogenic population collapse in orang-utans. *PLoS Biol* 4(2):285–291. doi: e2510.1371/journal.pbio.0040025
- Goudet J, Perrin N, Waser P (2002) Tests for sex-biased dispersal using bi-parentally inherited genetic markers. *Mol Ecol* 11(6):1103–1114
- Handley LJJ, Perrin N (2007) Advances in our understanding of mammalian sex-biased dispersal. *Mol Ecol* 16(8):1559–1578. doi:10.1111/j.1365-294X.2006.03152.x
- Hellborg L, Ellegren H (2003) Y chromosome conserved anchored tagged sequences (YCATS) for the analysis of mammalian male-specific DNA. *Mol Ecol* 12(1):283–291
- Li SZ, Wan HR, Ji HY, Zhou KY, Yang G (2009) SNP discovery based on CATS and genotyping in the finless porpoise

- (*Neophocaena phocaenoides*). *Conserv Genet* 10(6):2013–2019. doi:[10.1007/s10592-009-9882-4](https://doi.org/10.1007/s10592-009-9882-4)
- Morin PA, Chambers KE, Boesch C, Vigilant L (2001) Quantitative polymerase chain reaction analysis of DNA from noninvasive samples for accurate microsatellite genotyping of wild chimpanzees (*Pan troglodytes verus*). *Mol Ecol* 10(7):1835–1844
- Nater A, Krützen M, Lindholm AK (2008) Development of polymorphic microsatellite markers for the livebearing fish *Poecilia parac.* *Mol Ecol Resour* 8(4):857–860. doi:[10.1111/j.1755-0998.2008.02090.x](https://doi.org/10.1111/j.1755-0998.2008.02090.x)
- Nei M (1987) Molecular evolutionary genetics. Columbia University Press, New York
- Nietlisbach P (2009) Male-specific markers in orang-utans (*Pongo* spp.)—Dispersal and phylogeny. MSc thesis. University of Zurich
- Park SDE (2001) Trypanotolerance in West African cattle and the population genetic effects of selection. University of Dublin, Dublin, Ireland
- Reed DH, Frankham R (2003) Correlation between fitness and genetic diversity. *Conserv Biol* 17(1):230–237
- Rousset F (2008) GENEPOP '007: a complete re-implementation of the GENEPOP software for Windows and Linux. *Mol Ecol Resour* 8(1):103–106. doi:[10.1111/j.1471-8286.2007.01931.x](https://doi.org/10.1111/j.1471-8286.2007.01931.x)
- Utami SS, Goossens B, Bruford MW, de Ruiter JR, van Hooff J (2002) Male bimaturism and reproductive success in Sumatran orang-utans. *Behav Ecol* 13(5):643–652
- Warren KS, Verschoor EJ, Langenhuijzen S, Heriyanto, Swan RA, Vigilant L, Heeney JL (2001) Speciation and intrasubspecific variation of Bornean orang-utans, *Pongo pygmaeus pygmaeus*. *Mol Biol Evol* 18 (4):472–480
- Wich SA, Meijaard E, Marshall AJ, Husson S, Ancrenaz M, Lacy RC, van Schaik CP, Sugardjito J, Simorangkir T, Traylor-Holzer K, Doughty M, Supriatna J, Dennis R, Gumal M, Knott CD, Singleton I (2008) Distribution and conservation status of the orang-utan (*Pongo* spp.) on Borneo and Sumatra: how many remain? *Oryx* 42(3):329–339. doi:[10.1017/s003060530800197x](https://doi.org/10.1017/s003060530800197x)